



Elementary Complex Kleinian Groups

Mauricio Toledo-Acosta

Departamento de Matemáticas
Universidad de Sonora



Table of Contents

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

1 Introduction

2 Complex Kleinian Groups: An overview

3 Proof of the Theorem

4 Generalizations



Introduction

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Kleinian groups are discrete subgroups of $\mathrm{PSL}(2, \mathbb{C})$, the group of biholomorphic automorphisms of the complex projective line $\mathbb{CP}^1 \cong \mathbb{S}^2$, acting properly and discontinuously on a non-empty region of \mathbb{CP}^1 .



Introduction

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Kleinian groups are discrete subgroups of $\mathrm{PSL}(2, \mathbb{C})$, the group of biholomorphic automorphisms of the complex projective line $\mathbb{CP}^1 \cong \mathbb{S}^2$, acting properly and discontinuously on a non-empty region of \mathbb{CP}^1 .

Kleinian groups have been studied since the end of the 19th century by Fuchs, Klein, Poincaré, and many others. Kleinian groups have played a major role in several fields of mathematics, such as Riemann surfaces and Teichmüller theory, automorphic forms, holomorphic dynamics, conformal and hyperbolic geometry, etc.



Elementary Kleinian Groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The set of accumulation points of orbits of a Kleinian group is called the **limit set** of the group.



Elementary Kleinian Groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The set of accumulation points of orbits of a Kleinian group is called the **limit set** of the group.

Elementary groups are discrete subgroups of $\mathrm{PSL}(2, \mathbb{C})$ such that the limit set is a finite set. In other words, it is empty or it consists of 1 or 2 points.



Elementary Kleinian Groups

Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

The set of accumulation points of orbits of a Kleinian group is called the **limit set** of the group.

Elementary groups are discrete subgroups of $\text{PSL}(2, \mathbb{C})$ such that the limit set is a finite set. In other words, it is empty or it consists of 1 or 2 points.



(a) elliptic



(b) hyperbolic



(c) loxodromic



(d) parabolic



Generalize to higher dimensions

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Discrete subgroups of isometries of hyperbolic 3-space $\mathbb{H}_{\mathbb{R}}^3$ (discrete groups of conformal automorphisms of the sphere at infinity) \cong Discrete groups of holomorphic transformations of the complex projective line $\mathbb{CP}^1 \cong \mathbb{S}^2$ acting with nonempty region of discontinuity.

↓
Conformal Kleinian
Groups

↓
Complex Kleinian Groups



Table of Contents

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

1 Introduction

2 Complex Kleinian Groups: An overview

3 Proof of the Theorem

4 Generalizations



Preliminaries

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The **complex projective plane** \mathbb{CP}^2 is defined as

$$\mathbb{CP}^2 = (\mathbb{C}^3 \setminus \{0\}) / \mathbb{C}^*,$$

where \mathbb{C}^* acts by the usual scalar multiplication. Let

$$[\] : \mathbb{C}^3 \setminus \{0\} \rightarrow \mathbb{CP}$$

be the quotient map. We denote the projectivization of the point $x = (x_1, x_2, x_3) \in \mathbb{C}^3$ by $[x] = [x_1 : x_2 : x_3]$. We denote by e_1, e_2, e_3 the projectivization of the canonical base of \mathbb{C}^3 .



Preliminaries

Let $GL(3, \mathbb{C}) \subset \mathcal{M}_3(\mathbb{C})$ be the subgroup of matrices with determinant not equal to 0. The group of biholomorphic automorphisms of \mathbb{CP}^2 is given by

$$PSL(3, \mathbb{C}) := GL(3, \mathbb{C}) / \{\text{scalar matrices}\}.$$

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations



Preliminaries

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Let $GL(3, \mathbb{C}) \subset \mathcal{M}_3(\mathbb{C})$ be the subgroup of matrices with determinant not equal to 0. The group of biholomorphic automorphisms of \mathbb{CP}^2 is given by

$$PSL(3, \mathbb{C}) := GL(3, \mathbb{C}) / \{\text{scalar matrices}\}.$$

We denote the upper triangular subgroup of $PSL(3, \mathbb{C})$ by

$$U_+ = \left\{ \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{bmatrix} \mid a_{11}a_{22}a_{33} = 1, a_{ij} \in \mathbb{C} \right\}.$$



Preliminaries

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

As in the case of automorphisms of \mathbb{CP}^1 , we classify the elements of $\mathrm{PSL}(3, \mathbb{C})$ in three classes: elliptic, parabolic and loxodromic. However, unlike the classical case, there are several subclasses in each case. We now give a quick summary of the subclasses of elements we will be using.



Classification of elements of $\mathrm{PSL}(3, \mathbb{C})$

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

An element $g \in \mathrm{PSL}(3, \mathbb{C})$ is said to be:

- **Elliptic** if it has a diagonalizable lift in $\mathrm{SL}(3, \mathbb{C})$ such that every eigenvalue has norm 1.
- **Parabolic** if it has a non-diagonalizable lift in $\mathrm{SL}(3, \mathbb{C})$ such that every eigenvalue has norm 1.



Classification of elements of $\mathrm{PSL}(3, \mathbb{C})$

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- **Loxodromic** if it has a lift in $\mathrm{SL}(3, \mathbb{C})$ with an eigenvalue of norm distinct of 1. Furthermore, we say that g is:
 - **Loxo-parabolic**

$$\mathbf{h} = \begin{pmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda^{-2} \end{pmatrix}, |\lambda| \neq 1.$$

- A **complex homothety**, $\mathbf{h} = \mathrm{Diag}(\lambda, \lambda, \lambda^{-2})$, $|\lambda| \neq 1$.
- A **rational (resp. irrational) screw**, $\mathbf{h} = \mathrm{Diag}(\lambda_1, \lambda_2, \lambda_3)$, $|\lambda_1| = |\lambda_2| \neq |\lambda_3|$ and $\lambda_1 \lambda_2^{-1} = e^{2\pi i \theta}$ with $\theta \in \mathbb{Q}$ (resp. $\theta \in \mathbb{R} \setminus \mathbb{Q}$).
- **Strongly loxodromic**, $\mathbf{h} = \mathrm{Diag}(\lambda_1, \lambda_2, \lambda_3)$, where $\{|\lambda_1|, |\lambda_2|, |\lambda_3|\}$ are pairwise different.



Kernel of a Group

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Consider a subgroup $\Gamma \subset \mathrm{PSL}(3, \mathbb{C})$ acting on \mathbb{CP}^2 with a global fixed point $p \in \mathbb{CP}^2$. Let $\ell \subset \mathbb{CP}^2 \setminus \{p\}$ be a projective complex line. We define the projection $\pi = \pi_{p,\ell} : \mathbb{CP}^2 \rightarrow \ell$ given by $\pi(x) = \ell \cap \overleftrightarrow{px}$. This function is holomorphic, and it determines the group homomorphism

$$\Pi = \Pi_{p,\ell} : \mathrm{PSL}(3, \mathbb{C}) \rightarrow \mathrm{Bihol}(\ell) \cong \mathrm{PSL}(2, \mathbb{C})$$

given by $\Pi(g)(x) = \pi(g(x))$ for $g \in \Gamma$. We write $\mathrm{Ker}(\Gamma)$ instead of $\mathrm{Ker}(\Pi) \cap \Gamma$.

The **control group** of Γ is $\Pi(\Gamma)$.



Elementary Complex Kleinian Group

Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

How can we define *elementary* complex Kleinian groups?

- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines.



Elementary Complex Kleinian Group

Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

How can we define *elementary* complex Kleinian groups?

- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines.
- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines in general position.



Elementary Complex Kleinian Group

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

How can we define *elementary* complex Kleinian groups?

- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines.
- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines in general position.
- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ with reducible action.



Elementary Complex Kleinian Group

Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

How can we define *elementary* complex Kleinian groups?

- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines.
- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ such that its Kulkarni limit set contains a finite number of lines in general position.
- Discrete subgroups of $\mathrm{PSL}(3, \mathbb{C})$ with reducible action.
- Discrete solvable subgroups of $\mathrm{PSL}(3, \mathbb{C})$.



Solvable groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- If $g, h \in G$, we define the **commutator** as $[g, h] = g^{-1}h^{-1}gh$.
- We define the **commutator subgroup** as

$$[G, G] = \{[g, h] \mid g, h \in G\}.$$

- The **derived series** of G is given by

$$G^{(0)} = G, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}].$$

- We say that G is **solvable** if, for some $n \geq 0$, we have $G^{(n)} = \{id\}$.



Solvable groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- If $g, h \in G$, we define the **commutator** as $[g, h] = g^{-1}h^{-1}gh$.
- We define the **commutator subgroup** as

$$[G, G] = \{[g, h] \mid g, h \in G\}.$$

- The **derived series** of G is given by

$$G^{(0)} = G, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}].$$

- We say that G is **solvable** if, for some $n \geq 0$, we have $G^{(n)} = \{id\}$.



Solvable groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- If $g, h \in G$, we define the **commutator** as $[g, h] = g^{-1}h^{-1}gh$.
- We define the **commutator subgroup** as

$$[G, G] = \{[g, h] \mid g, h \in G\}.$$

- The **derived series** of G is given by

$$G^{(0)} = G, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}].$$

- We say that G is **solvable** if, for some $n \geq 0$, we have $G^{(n)} = \{id\}$.



Solvable groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- If $g, h \in G$, we define the **commutator** as $[g, h] = g^{-1}h^{-1}gh$.
- We define the **commutator subgroup** as

$$[G, G] = \{[g, h] \mid g, h \in G\}.$$

- The **derived series** of G is given by

$$G^{(0)} = G, \quad G^{(i+1)} = [G^{(i)}, G^{(i)}].$$

- We say that G is **solvable** if, for some $n \geq 0$, we have $G^{(n)} = \{id\}$.



Solvable groups: Examples

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- The **infinite dihedral group** is solvable

$$\mathrm{Dih}_\infty = \langle \mathrm{Rot}_\infty, z \mapsto -z \rangle.$$

- Any triangular group is solvable, with solvability length at most 3.
- The special orthogonal group is not solvable,

$$\left\{ \begin{bmatrix} a & -\bar{c} \\ c & \bar{a} \end{bmatrix} \mid |a|^2 + |b|^2 = 1 \right\} \subset \mathrm{PSL}(2, \mathbb{C})$$

- Cyclic groups, abelian groups.



Solvable groups: Examples

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- The **infinite dihedral group** is solvable

$$\mathrm{Dih}_\infty = \langle \mathrm{Rot}_\infty, z \mapsto -z \rangle.$$

- Any triangular group is solvable, with solvability length at most 3.
- The special orthogonal group is not solvable,

$$\left\{ \begin{bmatrix} a & -c \\ c & \bar{a} \end{bmatrix} \mid |a|^2 + |c|^2 = 1 \right\} \subset \mathrm{PSL}(2, \mathbb{C})$$

- Cyclic groups, abelian groups.



Solvable groups: Examples

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- The **infinite dihedral group** is solvable

$$\mathrm{Dih}_\infty = \langle \mathrm{Rot}_\infty, z \mapsto -z \rangle.$$

- Any triangular group is solvable, with solvability length at most 3.
- The special orthogonal group is not solvable,

$$\left\{ \begin{bmatrix} a & -c \\ c & \bar{a} \end{bmatrix} \mid |a|^2 + |c|^2 = 1 \right\} \subset \mathrm{PSL}(2, \mathbb{C})$$

- Cyclic groups, abelian groups.



Solvable groups: Examples

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

- The **infinite dihedral group** is solvable

$$\mathrm{Dih}_\infty = \langle \mathrm{Rot}_\infty, z \mapsto -z \rangle.$$

- Any triangular group is solvable, with solvability length at most 3.
- The special orthogonal group is not solvable,

$$\left\{ \begin{bmatrix} a & -c \\ c & \bar{a} \end{bmatrix} \mid |a|^2 + |b|^2 = 1 \right\} \subset \mathrm{PSL}(2, \mathbb{C})$$

- Cyclic groups, abelian groups.



The Kulkarni Limit Set

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

From now on, let $\Gamma \subset \mathrm{PSL}(3, \mathbb{C})$ be a discrete subgroup acting on \mathbb{CP}^2 .

Definition

- Let $L_0(\Gamma)$ be the closure of the set of points in \mathbb{CP}^n with infinite isotropy group.
- Let $L_1(\Gamma)$ be the closure of the set of cluster points of orbits of points in $\mathbb{CP}^n \setminus L_0(\Gamma)$.
- Let $L_2(\Gamma)$ be the closure of the set of cluster points of compact sets of $\mathbb{CP}^n \setminus (L_0(\Gamma) \cup L_1(\Gamma))$.

$$\Lambda_{Kul}(\Gamma) = \overline{L_0(\Gamma) \cup L_1(\Gamma) \cup L_2(\Gamma)}, \quad \Omega_{Kul}(\Gamma) = \mathbb{CP}^n \setminus \Lambda_{Kul}(\Gamma).$$



Limit sets for complex Kleinian groups

Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

The Kulkarni limit set $\Lambda_{\text{Kul}}(\Gamma)$ in \mathbb{CP}^2 is made up of points and complex projective lines. It contains 1, 2, 3, 4 or ∞ lines in general position.

The **equicontinuity region** for Γ , denoted $\text{Eq}(\Gamma)$, is defined to be the set of points $z \in \mathbb{CP}^n$ for which there is an open neighborhood U of z such that Γ restricted to U is a normal family. Γ acts properly and discontinuously on $\text{Eq}(\Gamma)$, and

$$\text{Eq}(\Gamma) \subset \Omega_{\text{Kul}}(\Gamma).$$



Limit sets for complex Kleinian groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The Kulkarni limit set $\Lambda_{\text{Kul}}(\Gamma)$ in \mathbb{CP}^2 is made up of points and complex projective lines. It contains 1, 2, 3, 4 or ∞ lines in general position.

The **equicontinuity region** for Γ , denoted $\text{Eq}(\Gamma)$, is defined to be the set of points $z \in \mathbb{CP}^n$ for which there is an open neighborhood U of z such that Γ restricted to U is a normal family.

Γ acts properly and discontinuously on $\text{Eq}(\Gamma)$, and

$$\text{Eq}(\Gamma) \subset \Omega_{\text{Kul}}(\Gamma).$$



Limit sets for complex Kleinian groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The Kulkarni limit set $\Lambda_{\text{Kul}}(\Gamma)$ in \mathbb{CP}^2 is made up of points and complex projective lines. It contains 1, 2, 3, 4 or ∞ lines in general position.

The **equicontinuity region** for Γ , denoted $\text{Eq}(\Gamma)$, is defined to be the set of points $z \in \mathbb{CP}^n$ for which there is an open neighborhood U of z such that Γ restricted to U is a normal family.

Γ acts properly and discontinuously on $\text{Eq}(\Gamma)$, and

$$\text{Eq}(\Gamma) \subset \Omega_{\text{Kul}}(\Gamma).$$



Limit sets for complex Kleinian groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The Kulkarni limit set $\Lambda_{\text{Kul}}(\Gamma)$ in \mathbb{CP}^2 is made up of points and complex projective lines. It contains 1, 2, 3, 4 or ∞ lines in general position.

The **equicontinuity region** for Γ , denoted $\text{Eq}(\Gamma)$, is defined to be the set of points $z \in \mathbb{CP}^n$ for which there is an open neighborhood U of z such that Γ restricted to U is a normal family. Γ acts properly and discontinuously on $\text{Eq}(\Gamma)$, and

$$\text{Eq}(\Gamma) \subset \Omega_{\text{Kul}}(\Gamma).$$



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible ✓



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible ✓

Reducible action {



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible ✓

Reducible action { Solvable



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible ✓

Reducible action $\begin{cases} \text{Solvable} \\ \text{Non-solvable} \end{cases}$



Background

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Action on \mathbb{CP}^2

Irreducible ✓

Reducible action $\left\{ \begin{array}{l} \text{Solvable} \\ \text{Non-solvable} \end{array} \right.$



Limit sets


Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations





Limit sets


Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations





Main Result

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset PSL(3, \mathbb{C})$ be a triangularizable complex Kleinian group such that its Kulkarni limit set does not consist of exactly four lines in general position. Then there exists a non-empty open region $\Omega_\Gamma \subset \mathbb{CP}^2$ such that

- (i) Ω_Γ is the maximal open set where the action is proper and discontinuous.*
- (ii) Ω_Γ is homeomorphic to one of the following regions: \mathbb{C}^2 , $\mathbb{C}^2 \setminus \{0\}$, $\mathbb{C} \times (\mathbb{H}^+ \cup \mathbb{H}^-)$ or $\mathbb{C} \times \mathbb{C}^*$.*
- (iii) Γ is finitely generated and $\text{rank}(\Gamma) \leq 4$.*



Main Result

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset PSL(3, \mathbb{C})$ be a triangularizable complex Kleinian group such that its Kulkarni limit set does not consist of exactly four lines in general position. Then

❖ *The group Γ can be written as*

$$\Gamma = \Gamma_p \rtimes \underbrace{\langle \eta_1 \rangle \rtimes \dots \rtimes \langle \eta_m \rangle}_{\text{loxo-parabolic}} \rtimes \underbrace{\langle \gamma_1 \rangle \rtimes \dots \rtimes \langle \gamma_n \rangle}_{\text{strongly loxodromic}}$$

where Γ_p is the subgroup of Γ consisting of all the parabolic elements of Γ .

❖ *The group Γ leaves a full flag invariant.*



Main Result

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset PSL(3, \mathbb{C})$ be a *solvable* complex Kleinian group such that its Kulkarni limit set does not consist of exactly four lines in general position. Then

❖ The group Γ can be written as

$$\Gamma = \Gamma_p \rtimes \underbrace{\langle \eta_1 \rangle \rtimes \dots \rtimes \langle \eta_m \rangle}_{\text{loxo-parabolic}} \rtimes \underbrace{\langle \gamma_1 \rangle \rtimes \dots \rtimes \langle \gamma_n \rangle}_{\text{strongly loxodromic}}$$

where Γ_p is the subgroup of Γ consisting of all the parabolic elements of Γ .

❖ The group Γ leaves a full flag invariant.



Table of Contents

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

1 Introduction

2 Complex Kleinian Groups: An overview

3 Proof of the Theorem

4 Generalizations



Ideas behind the proof

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The parabolic part is described in

Barrera, W., Cano, A., Navarrete, J. P., & Seade, J. (2022). Discrete parabolic groups in $\mathrm{PSL}(3, \mathbb{C})$. *Linear Algebra and its Applications*, 653, 430-500.



(v) Invariant Flag

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Borel fixed point theorem: Let G be a connected solvable group acting morphically on a non-empty complete variety V . Then G has a fixed point in V . Morphical action

Applying this theorem to the Zariski closure of Γ yields that Γ is virtually triangularizable. Namely, Γ has a finite index subgroup such that, up to conjugation, is upper triangular.

This proves (v).



(v) Invariant Flag

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Borel fixed point theorem: Let G be a connected solvable group acting morphically on a non-empty complete variety V . Then G has a fixed point in V . Morphical action

Applying this theorem to the Zariski closure of Γ yields that Γ is virtually triangularizable. Namely, Γ has a finite index subgroup such that, up to conjugation, is upper triangular.

This proves (v).



Conclusions (i)-(iv) are proved together.

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations



Restrictions on the Elements of a Non-commutative Group

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Proposition

Let $\Gamma \subset U_+$ be a discrete subgroup. Let $\gamma \in \Gamma$ be an irrational screw $\gamma = \text{Diag}(\beta^{-2}e^{-6\pi i\theta}, \beta e^{4\pi i\theta}, \beta e^{2\pi i\theta})$, for some $|\beta| \neq 1$ and $\theta \in \mathbb{R} \setminus \mathbb{Q}$, then Γ is commutative.

Proposition

Let $\Gamma \subset U_+$ be a non-commutative, torsion-free discrete subgroup, then Γ cannot contain a type I complex homothety.



The Core of a Group

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

The **core** of Γ is an important purely parabolic subgroup of a complex Kleinian group Γ which determines the dynamics of Γ .

Proposition

The elements of $\text{Core}(\Gamma)$ have the form

$$g_{x,y} = \begin{bmatrix} 1 & x & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

for some $x, y \in \mathbb{C}$.



The Core of a Group

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

It is straightforward to verify that

$$\Lambda_{\text{Kul}}(\text{Core}(\Gamma)) = \bigcup_{g_{x,y} \in \text{Core}(\Gamma)} \overleftrightarrow{e_1, [0 : -y : x]}$$

We denote this pencil of lines by $\mathcal{C}(\Gamma) = \Lambda_{\text{Kul}}(\text{Core}(\Gamma))$.

Proposition

Let $\Gamma \subset U_+$ be a discrete group, then every element of Γ leaves $\mathcal{C}(\Gamma)$ invariant.



Commutativity

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

$$\begin{cases} \Gamma \text{ is not commutative} \\ \Gamma \text{ is commutative} \end{cases}$$



Decomposition of Non-Commutative Triangular Groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset U_+$ be a non-commutative, torsion free, complex Kleinian group, then

$$\begin{aligned}\Gamma = \text{Core}(\Gamma) \rtimes \langle \xi_1 \rangle \rtimes \dots \rtimes \langle \xi_r \rangle \rtimes \\ \rtimes \langle \eta_1 \rangle \rtimes \dots \rtimes \langle \eta_m \rangle \rtimes \langle \gamma_1 \rangle \rtimes \dots \rtimes \langle \gamma_n \rangle.\end{aligned}$$

Furthermore, if $k = \text{rank}(\text{Core}(\Gamma))$ then $k + r + m + n \leq 4$.



Elementary Complex Kleinian Groups

Introduction

Complex Kleinian Groups: An overview

Proof of the Theorem

Generalizations

Parabolic

$$\begin{bmatrix} 1 & x & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{bmatrix}$$

$z \neq 0$

$\text{Core}(\Gamma)$

$A \setminus \text{Ker}(\Gamma)$

Loxodromic

$$\begin{bmatrix} \alpha & x & y \\ 0 & \beta & z \\ 0 & 0 & \beta \end{bmatrix} \quad \begin{bmatrix} \alpha & x & y \\ 0 & \beta & z \\ 0 & 0 & \gamma \end{bmatrix}$$

$\alpha \neq \beta, z \neq 0$ $\beta \neq \gamma$

Loxo-parabolic

Strongly loxodromic

$\text{Ker}(\lambda_{23}) \setminus A$



Morphisms λ

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Let $\lambda_{12}, \lambda_{23}, \lambda_{13} : (U_+, \cdot) \rightarrow (\mathbb{C}^*, \cdot)$ be the group morphisms given by

$$\begin{aligned}\lambda_{12}([\alpha_{ij}]) &= \alpha_{11}\alpha_{22}^{-1} \\ \lambda_{23}([\alpha_{ij}]) &= \alpha_{22}\alpha_{33}^{-1} \\ \lambda_{13}([\alpha_{ij}]) &= \alpha_{11}\alpha_{33}^{-1}.\end{aligned}$$

Strategy of the proof:

- Decomposition of Γ in terms of $\text{Ker}(\lambda_{23})$.
- Decompose $\text{Ker}(\lambda_{23})$ in terms of $\text{Ker}(\lambda_{12})$.
- Decompose $A = \text{Ker}(\lambda_{12}) \cap \text{Ker}(\lambda_{23})$ in terms of $\text{Ker}(\Gamma)$



Rank

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem (Bestvina, Kapovich, Kleiner)

Let Γ be a group acting properly and discontinuously on a contractible manifold of dimension m , then $\text{obdim}(\Gamma) \leq m$.



Rank

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem (Bestvina, Kapovich, Kleiner)

Let Γ be a group acting properly and discontinuously on a contractible manifold of dimension m , then $\text{obdim}(\Gamma) \leq m$.

In our case, it can be re-stated as:

Theorem

Let $\Gamma \subset U_+$ be a non-commutative, torsion free, complex Kleinian group acting properly and discontinuously on a simply connected domain $\Omega \subset \mathbb{CP}^2$, then $k + r + m + n \leq 4$.



Rank

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem (Bestvina, Kapovich, Kleiner)

Let Γ be a group acting properly and discontinuously on a contractible manifold of dimension m , then $\text{obdim}(\Gamma) \leq m$.

In our case, it can be re-stated as:

Theorem

Let $\Gamma \subset U_+$ be a non-commutative, torsion free, complex Kleinian group acting properly and discontinuously on a simply connected domain $\Omega \subset \mathbb{CP}^2$, then $k + r + m + n \leq 4$.

Find a simply connected domain $\Omega \subset \mathbb{CP}^2$ where Γ acts properly and discontinuously, and then apply the theorem. In some cases, we write the explicit decomposition of Γ and verify that $\text{rank}(\Gamma) \leq 4$.



Some cases

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Denote $\Sigma = \Pi(\Gamma)$. If Σ is discrete and $\text{Ker}(\Gamma)$ is finite. If $|\Lambda(\Sigma)| \neq 2$, let

$$\Omega = \left(\bigcup_{z \in \Omega(\Sigma)} \overleftrightarrow{e_1, z} \right) \setminus \{e_1\}.$$

We know that Γ acts properly and discontinuously on Ω . If $|\Lambda(\Sigma)| = 0, 1$ or ∞ , then each connected component of Ω is simply connected, since they are respectively homeomorphic to \mathbb{CP}^2 , \mathbb{C}^2 or $\mathbb{C} \times \mathbb{H}$. By the theorem, it follows $k + r + m + n \leq 4$



Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

For non-commutative Γ , using these ideas, we have constructed an open subset $\Omega_\Gamma \subset \mathbb{CP}^2$ such that the orbits of every compact set $K \subset \Omega_\Gamma$ accumulate on $\mathbb{CP}^2 \setminus \Omega_\Gamma$. Thus we can define a limit set for the action of Γ by $\Lambda_\Gamma := \mathbb{CP}^2 \setminus \Omega_\Gamma$. This limit set describes the dynamics of Γ , and the open region Ω_Γ satisfies (i) and (ii).

Also, we prove that $\text{rank}(\Gamma) \leq 4$. This verifies (iii).



Commutative groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem (Barrera, Cano, Navarrete, Seade)

Let $\Gamma \subset U_+$ be a commutative group, then Γ is conjugate in $PSL(3, \mathbb{C})$ to a subgroup of one of the following Abelian Lie Groups:



$$C_1 = \left\{ \begin{pmatrix} \alpha^{-2} & 0 & 0 \\ 0 & \alpha & \beta \\ 0 & 0 & \alpha \end{pmatrix} \mid \alpha \in \mathbb{C}^*, \beta \in \mathbb{C} \right\}.$$



$$C_2 = \{ \text{Diag}(\alpha, \beta, \alpha^{-1}\beta^{-1}) \mid \alpha, \beta \in \mathbb{C}^* \}.$$



$$C_3 = \left\{ \begin{pmatrix} 1 & 0 & \beta \\ 0 & 1 & \gamma \\ 0 & 0 & 1 \end{pmatrix} \mid \beta, \gamma \in \mathbb{C} \right\}.$$



Commutative groups

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem (Barrera, Cano, Navarrete, Seade)

•

$$C_4 = \left\{ \begin{pmatrix} 1 & \beta & \gamma \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \mid \beta, \gamma \in \mathbb{C} \right\}.$$

•

$$C_5 = \left\{ \begin{pmatrix} 1 & \beta & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix} \mid \beta, \gamma \in \mathbb{C} \right\}.$$



Case 1: Form

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Proposition

Let $\Gamma \subset U_+$ be a commutative subgroup such that each element of Γ has the form C_1 . Then there exists an additive subgroup $W \subset (\mathbb{C}, +)$, and a group morphism $\mu : (W, +) \rightarrow (\mathbb{C}^, \cdot)$ such that*

$$\Gamma = \Gamma_{W, \mu} = \left\{ \begin{bmatrix} \mu(w)^{-2} & 0 & 0 \\ 0 & \mu(w) & w\mu(w) \\ 0 & 0 & \mu(w) \end{bmatrix} \mid w \in W \right\}.$$



Case 1: Discreteness and Rank

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Proposition

Let $\Gamma = \Gamma_{W,\mu} \subset U_+$ be a group as described in previous proposition. Γ is discrete if and only if $\text{rank}(W) \leq 3$ and the morphism μ satisfies the following condition:

- *Whenever we have a sequence $\{w_k\} \in W$ of distinct elements such that $w_k \rightarrow 0$, either $\mu(w_k) \rightarrow 0$ or $\mu(w_k) \rightarrow \infty$.*



Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Case	Conditions
C1.1	$\mu(W)$ has rational rotations and W is discrete.
C1.2	$\mu(W)$ has rational rotations and W is not discrete.
C1.3	$\mu(W)$ has no rational rotations but has irrational rotations, and W is discrete.
C1.4	$\mu(W)$ has no rational or irrational rotations, and W is discrete.
C1.5	$\mu(W)$ has no rational rotations but has irrational rotations, and W is not discrete.
C1.6	$\mu(W)$ has no rational or irrational rotations, and W is not discrete.



Case 1: Kulkarni Limit Set

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset PSL(3, \mathbb{C})$ be a commutative discrete group having the form given previous proposition, then

$$\Lambda_{Kul}(\Gamma) = \begin{cases} \overleftrightarrow{e_1, e_2}, & \begin{cases} \text{Cases C1.3 or C1.4, with condition} \\ \text{Case C1.1} \end{cases} \\ \{e_1\} \cup \overleftrightarrow{e_2, e_3}, & \text{Cases C1.5 or C1.6 no condition (F).} \\ \overleftrightarrow{e_1, e_2} \cup \overleftrightarrow{e_2, e_3}, & \begin{cases} \text{Cases C1.5 or C1.6, with condition} \\ \text{Case C1.2} \end{cases} \end{cases}$$



Case 2: Form

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Proposition

Let $\Gamma \subset U_+$ be a commutative subgroup such that each element of Γ has the form $\text{Diag}(\alpha, \beta, \alpha^{-1}\beta^{-1})$. Then there exist two multiplicative subgroups $W_1, W_2 \subset (\mathbb{C}^, \cdot)$ such that*

$$\Gamma = \Gamma_{W_1, W_2} = \{ \text{Diag}(w_1, w_2, 1) \mid w_1 \in W_1, w_2 \in W_2 \}.$$



Case 2: Rank

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Proposition

Let $\Gamma \subset U_+$ be a diagonal discrete group such that every element has the form $\gamma = \text{Diag}(w_1, w_2, 1)$. Then $\text{rank}(\Gamma) \leq 2$.



Case 2

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

If $\alpha^n = \beta^m$ for some $n, m \in \mathbb{Z}$:

[D1] $L_0(\Gamma) \cup L_1(\Gamma) = \overleftrightarrow{e_1, e_2} \cup \{e_3\}$, if $|\alpha| > 1 > |\beta|$ or $|\alpha| < 1 < |\beta|$.

[D2] $L_0(\Gamma) \cup L_1(\Gamma) = \overleftrightarrow{e_1, e_2} \cup \{e_3\}$, if $|\alpha| > |\beta| > 1$ or $|\alpha| < |\beta| < 1$.

If there are no integers n, m such that $\alpha^n = \beta^m$:

[D3] $L_0(\Gamma) \cup L_1(\Gamma) = \{e_1, e_2, e_3\}$, if $|\alpha| > 1 > |\beta|$ or $|\alpha| < 1 < |\beta|$.

[D4] $L_0(\Gamma) \cup L_1(\Gamma) = \{e_1, e_2, e_3\}$, if $|\alpha| > |\beta| > 1$ or $|\alpha| < |\beta| < 1$.

[D5] $L_0(\Gamma) \cup L_1(\Gamma) = \overleftrightarrow{e_1, e_2} \cup \overleftrightarrow{e_2, e_3}$, if β is an irrational rotation.



Case 2: Kulkarni Limit Set

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma_{\alpha,\beta} \subset U_+$ be a discrete group containing loxodromic elements, then

- i) $\Lambda_{Kul}(\Gamma) = \overleftrightarrow{e_1, e_2} \cup \{e_3\}$ in Cases [D1] and [D2].
- ii) $\Lambda_{Kul}(\Gamma) = \{e_1, e_2, e_3\}$ in Cases [D3] and [D4].
- iii) $\Lambda_{Kul}(\Gamma) = \overleftrightarrow{e_1, e_2} \cup \overleftrightarrow{e_2, e_3}$ in Case [D5].



Commutative case: Proof of the Main Theorem

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

If Γ is commutative, it is conjugate to a subgroup of the Lie groups C_1 or C_2 . In this setting, the region $\Omega_{\text{Kul}}(\Gamma)$ satisfies conclusions (i) and (ii) as a consequence of the previous theorems. Again, $\text{rank}(\Gamma) \leq 4$, this proves conclusion (iii).

On the other hand, $\Gamma \cong \mathbb{Z}^r$ with $r = \text{rank}(\Gamma)$, and then we can write Γ as a trivial semidirect product of copies of \mathbb{Z} , thus verifying conclusion (iv).



Table of Contents

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

1 Introduction

2 Complex Kleinian Groups: An overview

3 Proof of the Theorem

4 Generalizations



A First Generalization ✓

Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Theorem

Let $\Gamma \subset \mathrm{PSL}(3, \mathbb{C})$ be a solvable complex Kleinian group such that its Kulkarni limit set does not consist of exactly four lines in general position. Let $\Gamma_0 \subset \Gamma$ be a virtually triangularizable finite index subgroup. If Γ_0 is commutative then there exists a non-empty open region $\Omega_\Gamma \subset \mathbb{CP}^2$ such that

- (i) Ω_Γ is the maximal open set where the action is proper and discontinuous.*
- (ii) Ω_Γ is homeomorphic to one of the following regions: \mathbb{C}^2 , $\mathbb{C}^2 \setminus \{0\}$, $\mathbb{C} \times (\mathbb{H}^+ \cup \mathbb{H}^-)$ or $\mathbb{C} \times \mathbb{C}^*$.*
- (iii) Up to a finite index subgroup, the group Γ leaves a full flag invariant.*



Elementary
Complex
Kleinian
Groups

Introduction

Complex
Kleinian
Groups: An
overview

Proof of the
Theorem

Generalizations

Thank you



Table of Contents

Elementary
Complex
Kleinian
Groups

Appendix

5 Appendix



A Group Acting Morphically

Elementary
Complex
Kleinian
Groups

Appendix

Definition

Let G be an algebraic group, V a variety, and let $\alpha : G \times V \rightarrow V$ be an action of the group G in V , $(g, x) \mapsto gx = \alpha(g, x)$. One says that G *acts morphically* on V if the action α satisfies the following axioms:

- ❶ $\alpha(e, x) = x$, for any $x \in V$, where $e \in G$ is the identity element.
- ❷ $\alpha(g, hx) = \alpha(gh, x)$ for any $g, h \in G$ and $x \in V$.

Solvable groups are virtually triangular